

This is a repository copy of *Latitudinal constraints in responsiveness of plants to arbuscular mycorrhiza : the 'sun-worshipper' hypothesis*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/146047/>

Version: Accepted Version

Article:

Veresoglou, Stavros, Chen, Baodong, Fischer, Matthias et al. (5 more authors) (2019) Latitudinal constraints in responsiveness of plants to arbuscular mycorrhiza : the 'sun-worshipper' hypothesis. *New Phytologist*. ISSN 1469-8137

<https://doi.org/10.1111/nph.15918>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



New Phytologist

Latitudinal constraints in responsiveness of plants to arbuscular mycorrhiza: The 'sun-worshipper' hypothesis.

Journal:	<i>New Phytologist</i>
Manuscript ID	NPH-L-2019-29648.R1
Manuscript Type:	L - Letter
Date Submitted by the Author:	n/a
Complete List of Authors:	Veresoglou, Stavros; Freie Universitat Berlin, Plant Ecology Chen, Baodong; Research Centre for Eco-Environmental Sciences, Environmental Sciences Fischer, Matthias; Freie Universitat Berlin, Plant Ecology Helgason, Thorunn; University of York, Department of Biology Mamolos, Andreas; Aristotle University of Thessaloniki, Faculty of Agriculture Rillig, Matthias; Freie Universitat Berlin Fachbereich Biologie Chemie Pharmazie, Institut für Biologie Roldán, Antonio; CSIC, Department of Soil and Water Conservation Johnson, David; University of Manchester, School of Earth and Environmental Sciences
Key Words:	Arbuscular mycorrhiza, eco-evolutionary processes, environmental filtering, Glomeromycotina, mycorrhizal responsiveness, phenotypic adaptation, plant biogeography, the sun-worshipper hypothesis

SCHOLARONE™
Manuscripts

Letter

Number of Words (summary): 0
Number of Words (main text): 1496
Number of references: 31
Display items: 2 figures no tables, no text boxes

Latitudinal constraints in responsiveness of plants to arbuscular mycorrhiza: The ‘sun-worshipper’ hypothesis.

Stavros D. Veresoglou^{1,2*}, Baodong Chen^{3,4}, Matthias M. Fischer¹, Thorunn Helgason⁵,
Andreas P. Mamolos², Matthias C. Rillig^{1,6}, Antonio Roldán⁷, David Johnson⁸

¹ Freie Universität Berlin, Institut für Biologie, Plant Ecology, D-14195 Berlin, Germany

² Faculty of Agriculture, Laboratory of Ecology and Environmental Protection, Aristotle University of Thessaloniki, 541 24 Thessaloniki, Greece.

³ State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, China

⁴ University of Chinese Academy of Sciences, Beijing, China

⁵ Department of Biology, University of York, York, YO10 5DD, UK

⁶ Berlin-Brandenburg Institute of Advanced Biodiversity Research (BBIB), D-14195 Berlin, Germany

⁷ CSIC-Centro de Edafología y Biología Aplicada del Segura, Department of Soil and Water Conservation, P.O. Box 164, Campus de Espinardo, 30100 Murcia, Spain

⁸ School of Earth and Environmental Sciences, The University of Manchester, Manchester, M13 9PT. UK.

* Email address: sveresoglou@zedat.fu-berlin.de, telephone: +49 30 838-53172

Keywords

Arbuscular mycorrhiza; eco-evolutionary processes; environmental filtering;
Glomeromycotina; mycorrhizal responsiveness; phenotypic adaptation; plant biogeography;
the sun-worshipper hypothesis

Most terrestrial plants depend strongly on associations with arbuscular mycorrhizal (AM) fungi (Subphylum: Glomeromycotina) to establish and survive (Bever, 2002; van der Heijden *et al.*, 1998; Klironomos *et al.*, 2011; Veresoglou *et al.*, 2017), and have evolved a nutritional mutualism. In this mutualism, the plant provides carbon to the fungus, usually subject to the availability of light (Hayman 1970; Heinemeyer *et al.*, 2003; Shi *et al.* 2014; Konvalinkova & Jansa, 2016), and the fungus provides the plant with mineral nutrients acquired from soil. Because of light constraints, it is expected that latitude exerts a strong influence on reciprocal exchange of resources between mycorrhizal plants and fungi, and this could have consequences on the responsiveness of plants to mycorrhizal fungi. Latitude induces changes in the amount of solar energy and the timing when this is made available to primary producers during the year and in the day. At the same time, there is a strong negative relationship between latitude and temperature that may also impact the functioning of the mycorrhizal symbiosis, and in some cases (e.g. in north temperate systems), a general relationship between latitude and several edaphic factors (Read and Perez-Moreno 2003).

There is compelling evidence that the alpha-diversity of Glomeromycotan fungi, which form AM symbioses, decreases with latitude (Davison *et al.*, 2015). This finding can be partially explained by the transition from ecosystems dominated by AM host plants in the tropics, to ectomycorrhizal and ericoid mycorrhizal dominated ecosystems at higher latitudes (Smith & Read, 2008). We know less about the extent to which latitude impacts the functioning of AM symbioses, which could occur either through latitudinal differences in solar radiation or resulting changes in temperature (Clarke & Gaston, 2006; Schluter, 2016). Here, we propose the ‘sun-worshipper’ hypothesis that discriminates three different types of underlying responses of latitudinal gradient-induced changes in plant host mycorrhizal responsiveness (Fig. 1). Changes in abiotic conditions may allow plants to derive more benefits from the symbiosis at lower latitudes through phenotypic plasticity (mechanism one – Fig.1). A likely example of phenotypic plasticity might involve changes in the expression of genes that allow crosstalk with AM fungi when light availability is low as has been shown for drought (Li *et al.*, 2016). Abiotic conditions more favourable for the symbiosis close to the tropics could further exclude, via competition, species less dependent on AM fungi, resulting in distinct plant communities from a perspective of AM fungal-associating behaviour; such a process can be described as environmental filtering (mechanism two – Fig. 1). Finally, we know that AM plants at high latitudes encounter a less diverse pool of potential symbiotic partners (e.g. because of the observed latitudinal gradient in Glomeromycotan diversity; Davison *et al.*, 2015) and at the same time communities at high latitudes are in general

dominated by plants that associate with ectomycorrhizal and ericoid mycorrhizal fungi. Plants distant from the tropics could thus form less profitable AM symbioses (but also support fewer AM partners) because of a more limited pool of suitable AM fungal partners, and this mechanism is analogous to the indirect eco-evolutionary causes (Pärtel, 2002) (mechanism three - Fig. 1). Here, we use the term eco-evolutionary processes to describe combined effects of latitude on phenotypic plasticity, environmental filtering and eco-evolutionary adaptation of the host plant trait mycorrhizal dependency (e.g. Thuiller *et al.*, 2013). Even though these different mechanisms are not mutually exclusive, it is important to disentangle how each of them influences how host plants respond to mycorrhizal symbioses along gradients of latitude and solar radiation.

The benefits that plants receive from the symbiosis in relation to the carbon costs vary considerably depending on abiotic growth conditions (Johnson *et al.*, 1997; Hoeksema *et al.* 2010; Grman and Robinson, 2013), compatibility of the plant host with the local AM fungal community (Klironomos, 2003) and the degree to which a plant can take advantage of non-nutritional functions of mycorrhiza such as protection from pathogens (Veresoglou & Rillig, 2013). Resource stoichiometry of phosphorus (P), nitrogen (N) and light, in particular, represents a proven tool explaining variance in growth responses of mycorrhizal hosts at various spatial scales (Johnson, 2010). Latitude-related predictions could complement such existing tools in understanding why mycorrhizal growth responses differ at large scales. It may additionally illuminate systematic differences in mycorrhizal responsiveness such as those explained by the life history of the hosts (Boerner, 1992; Roumet *et al.*, 2005). This would be the case if the latitudinal effects are mediated through differences in solar radiation. Annual terrestrial plants may never experience light-duration stress during winter, whereas the opposite is the case for perennials that represent the majority of terrestrial plants (for example over 70% of species in the LEDA database are perennials – Kleyer *et al.* 2008). There is good evidence that perennial AM fungi can survive over winter in the roots of their hosts (Buwalda *et al.*, 1985; Dodd & Jeffries, 1986 but see Hetrick *et al.*, 1984; Mohammed *et al.*, 1988), which could affect the carbon economy of their plant hosts. During winter, plant requirements for nutrients are limited and photoassimilates are in short supply; therefore, plants that can confine the activity of their mycorrhizal partners may benefit through improved survival rates. As a result, we expect that there is evolutionary pressure for perennials to further confine mycorrhizal responsiveness when growing outside the tropics, compared to annuals.

We undertook three synthesis activities to establish whether the expectations outlined above are plausible for mycorrhizal systems (see Methods in supplementary material). We first compiled a database on crop plant responses to mycorrhiza to identify phenotypic responses to latitude (Fig. S1). To assess environmental filtering due to AM fungi with regards to latitude, we synthesized data from a common garden experiment on comparative mycorrhizal responsiveness of North-American annual and perennial herbaceous plants (Wilson & Hartnett, 1998) with plant distribution data for the specific plants from USDA (USDA, 2016). We also tested for differences in mycorrhizal responsiveness across genotypes of *Zea mays* (maize) that are routinely used either in temperate or tropical systems, despite that genetic variability could effectively be attributed to breeding. These syntheses activities were not sufficiently robust to address the mechanistic constituents of the sun worshipper hypothesis but were carried out to support the over-arching concept and encourage larger syntheses or experiments exploring the hypothesis in the future.

Variance in the database on crop plant responses could be best explained, in our models consisting of a single predictor, by photosynthetic radiation (Fig 2.a). The optimal model had an intercept of -0.5 ($F=9.7$, $P<0.001$) and a slope per $\text{MJ.m}^{-2}.\text{day}^{-1}$ radiation of 0.23 ($F=6.1$, $P=0.019$). Fitted intercepts for the different plant species shared a standard deviation of 0.048 (Fig. 2a). Analysis of maize lines demonstrated that eco-evolutionary processes also drive latitude-dependencies on mycorrhizal responsiveness. The Mann-Whitney test between temperate- (i.e. middle two quartiles) and tropical-climate adapted lines of maize revealed higher responsiveness for temperate lines ($U = 24.5$, $P=0.034$). When we repeated this analysis for tropical lines through maintaining the two middle quartiles, the differences became even more apparent ($U = 4$, $P<0.001$, Fig. 2b). Mycorrhizal responsiveness of species in Wilson and Hartnett (1998) could be predicted by latitude of their distribution, which suggests that mycorrhizal responsiveness might induce an environmental filtering. We raised latitude to the fourth power to address fitting issues and obtained an intercept of 0.65; latitude slope of $-2.44 \cdot 10^{-8}$ ($F_{\text{lat}}=23.32$; $P<0.001$; $R^2_{\text{adj}}=0.19$; Fig. 2c). We subsequently fitted an additional parameter that differentiated between annual and perennial plants. Inclusion of the categorical variable perennial was significant ($F=23.36$, $P<0.001$) and there was a significant interaction between this parameter and latitude ($F=9.47$, $P=0.003$) suggesting that slopes also differed. When we analyzed annual and perennial plants separately, we found a significant relationship only for perennials (Kendall Tau was -0.32 - $P=0.004$, whereas for annuals the Kendall Tau was 0.03 - $P=0.87$).

The sun-worshipper hypothesis predicts that latitude impacts mycorrhizal responsiveness in plants via three complementary mechanisms, namely phenotypic responses, eco-evolutionary processes and environmental filtering (Fig. 1). Even though the analyses we report have limited resolution, they were supportive of the sun worshipper hypothesis. We found evidence that phenotypic plasticity (Fig. 2a), eco-evolutionary processes (Fig. 2b) and environmental filtering (Fig. 2c) might be operational for all three different types of latitude related differences in mycorrhizal responsiveness (Thuiller *et al.*, 2013). Despite the findings from our analyses, it is important to highlight additional factors that may also influence mycorrhizal responsiveness, and which have the potential to confound our findings. For example, our observations may correlate with systematic differences in soil fertility; high weathering rates generally lead to poorer fertility, as occurs in many parts of the tropics (Read and Perez-Moreno, 2003). Therefore, disentangling the specific role of light versus other edaphic and environmental factors in driving mycorrhizal responsiveness likely requires additional experimentation. Testing competing hypotheses could be done with carefully designed common garden experiments or synthesizing evidence from altitudinal experiments. Nevertheless, our analysis and associated hypothesis prompts further mechanistic analyses to test how resource stoichiometry and other critical functions undertaken by mycorrhizal fungi are influenced by latitudinal gradients.

Acknowledgements

SDV is partly supported by the DFG project Metacorrhiza (VE 736/2-1). DJ is partly supported by the N8 AgriFood programme and NERC. TH is partly supported by the BBSRC grant BB/L026007/1. The letter has benefited considerably from comments and suggestions from Ian Dickie and three anonymous reviewers.

Contributions

SDV: Conceived the project and carried out the analyses; SDV TH, APM, MCR, AR, DJ: Discussed and further developed the idea; SDV, DJ: Wrote the manuscript; All authors commented to the manuscript and approved the final version of it.

Supplementary Information

Appendix 1: Mycorrhizal dependency as proxy of investment

Appendix 2: Meta-analysis on phenotypic plasticity

Appendix 3: Eco-evolutionary processes on LGMR

Appendix 4: Environmental filtering on LGMR

Fig. S1: Locations of the 40 studies included in the meta-analysis

Table S1. Primary data on the 42 studies which we considered in our meta-analysis.

References

- Bever JD. 2002.** Negative feedback within a mutualism: host-specific growth of mycorrhizal fungi reduces plant benefit. *Proceedings of the Royal Society B* **269**, 2595-2601.
- Boerner REJ. 1992.** Plant life span and response to inoculation with vesicular-arbuscular mycorrhizal fungi. *Mycorrhiza* **1**, 153-161.
- Buwalda JG, Stribley DP, Tinker PB. 1985.** VA-mycorrhiza of winter and spring cereals. *Journal of Agricultural Science* **105**, 649-657.
- Clarke A, Gaston KJ. 2006.** Climate, energy and diversity. *Proceedings of the Royal Society B* **273**, 2257-2266.
- Davison J, Moora M, Öpik M, Adholeya A, Ainsaar L, Bâ A, et al. 2015.** Global assessment of arbuscular mycorrhizal fungus diversity reveals very low endemism. *Science* **349**: 970-973.
- Dodd JC, Jeffries P. 1986.** Early development of VAM in autumn-sown cereals. *Soil Biology and Biochemistry* **18**, 149-154
- Grman E, Robinson TMP. 2013.** Resource availability and imbalance affect plant–mycorrhizal interactions: a field test of three hypotheses. *Ecology* **94**, 62-71.
- Hayman DS. 1974.** Plant growth responses to vesicular-arbuscular mycorrhiza .6 effect of light and temperature. *New Phytologist* **73**, 71-&
- Heinemeyer A, Ringway KP, Edwards EJ, Benham DG, Young PW, Fitter AH. 2003.** Impact of soil warming and shading on colonization and community structure of arbuscular mycorrhizal fungi in roots of a native grassland community. *Global Change Biology* **10**, 52-64.
- Hetrick BAD, Bockus WW, Bloom J. 1984.** The role of vesicular-arbuscular mycorrhizal fungi in the growth of Kansas winter wheat. *Canadian Journal of Botany* **62**, 735-740.
- Hoeksema JD, Chaudhary VB, Gehring CA, Johnson NC, Karst J, Koide RT, Pringle A, Zabiski C, Bever JD, Moore JC. et al. 2010.** A meta-analysis of context-dependency in plant response to inoculation with mycorrhizal fungi. *Ecology Letters* **13**, 394-407.
- Johnson NC, Graham JH, Smith FA. 1997.** Functioning of mycorrhizal associations along the mutualism-parasitism continuum. *New Phytologist* **135**, 575–585.

- 204 **Johnson NC. 2010.** Resource stoichiometry elucidates the structure and function of
 205 arbuscular mycorrhizas across scales. *New Phytologist* **185**, 631-647.
- 206 **Kleyer M, Bekker R M, Knevel I C, Bakker J P, Thompson K, Sonnenschein M and et al.**
 207 **2008.** The LEDA Traitbase: A database of life-history traits of Northwest European
 208 flora. *Journal of Ecology* **96**, 1266-1274.
- 209 **Klironomos JN, Zobel M, Tibbett M, Stock WD, Rillig MC, Parrent JL, Moora M, Koch**
 210 **AM, Facelli JM, Facelli E, Dickie IA, Bever JD. 2011.** Forces that structure plant
 211 communities: quantifying the importance of the mycorrhizal symbiosis. *New*
 212 *Phytologist* **189**, 366-370.
- 213 **Konvalinkova T, Jansa J. 2016.** Lights off for arbuscular mycorrhiza: On its symbiotic
 214 functioning under light deprivation. *Frontiers in Plant Science* **7**, 782.
- 215 **Li T, Sun Y, Ruan Y, Xu L, Hu Y, Hao Z, Zhang X, Li H, Wang Y, Yang L, Chen B.**
 216 **2016.** Potential role of D-myo-inositol-3-phosphate synthase and 14-3-3 genes in the
 217 crosstalk between *Zea mays* and *Rhizophagus intraradices* under drought stress.
 218 *Mycorrhiza* **26**, 879-893.
- 219 **Mohammad MJ, Pan WL, Kennedy AC. 1998.** Seasonal mycorrhizal colonization of winter
 220 wheat and its effect on wheat growth under dryland field conditions. *Mycorrhiza* **8**,
 221 139-144.
- 222 **Pärtel M. 2002.** Local plant diversity patterns and evolutionary history at the regional scale.
 223 *Ecology* **83**, 2361-2366.
- 224 **Read DJ, Perez-Moreno J. 2003.** Mycorrhizas and nutrient cycling in ecosystems – a
 225 journey towards relevance? *New Phytologist* **157**, 475-492.
- 226 **Roumet C, Urcelay C, Díaz S. 2005.** Suites of root traits differ between annual and perennial
 227 species growing in the field. *New Phytologist* **170**, 357-368.
- 228 **Schluter D. 2016.** Speciation, Ecological Opportunity, and Latitude. *American Naturalist* **187**,
 229 1-18.
- 230 **Shi G, Liu Y, Johnson NC, Olsson PA, Mao L, Cheng G, Jiang S, An L, Du G, Feng H.**
 231 **2014.** Interactive influence of light intensity and soil fertility on root-associated
 232 arbuscular mycorrhizal fungi. *Plant and Soil* **378**, 173-188.
- 233 **Smith SE, Read DJ. 2008.** Mycorrhizal symbiosis, 33d ed. Academic Press, London, UK.
- 234 **Tedersoo L, Nara K. 2010.** General latitudinal gradient of biodiversity is reversed in
 235 ectomycorrhizal fungi. *New Phytologist* **185**, 351-354.

- Thuiller W, Münkemüller T, Lavergne S, Mouillot S, Mousquet N, Schiffrers K, Gravel D. 2013.** A road map for integrating eco-evolutionary processes into biodiversity models. *Ecology Letters* **16**, 94-105.
- USDA, NRCS. 2016.** The PLANTS Database (<http://plants.usda.gov>, 28 January 2017). National Plant Data Team, Greensboro, NC 27401-4901 USA.
- van der Heijden MGA, Klironomos JN, Ursic M, Moutoglis P, Streitwolf-Engel R, Boller T, Wiemken A, Sanders IR. 1998.** Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature* **396**, 72-75.
- Veresoglou SD, Rillig MC. 2013** Accounting for the adaptation deficit of non-mycorrhizal plants in experiments. *Plant and Soil* **366**, 33-34.
- Veresoglou SD, Wulf M, Rillig MC. 2017.** Facilitation between woody and herbaceous plants that associate with arbuscular mycorrhizal fungi in temperate European forests. *Ecology and Evolution* **7**, 1181-1189.
- Wilson GWT, Hartnett DC. 1998.** Interspecific variation in plant responses to mycorrhizal colonization in tallgrass prairie. *American Journal of Botany* **85**, 1732-1738.

Legends to Figures

Fig. 1. The ‘sun-worshipper’ hypothesis: Light availability declines with latitude (x -axis) to which plant species that associate with AM fungi respond via reducing their mycorrhizal responsiveness (LGMR – y -axis right). This is depicted in the figure with the black continuous line describing phenotypic plasticity (Mechanism 1). Eco-evolutionary responses partially offset the decline in LGMR with latitude (the resulting black discontinuous line has a lower slope - Mechanism 2). Reduced mycorrhizal responsiveness further induces a decline in species that associate with AM fungi (red line) with species that either associate with ectomycorrhizal (ECM) or ericoid (ERM) mycorrhizal fungi (green continuous line – y -axis left). This represents Mechanism 3. As a result, latitude is a key factor that determining plant responsiveness to mycorrhizal fungi

Fig. 2 (a) Scatterplot of mycorrhizal responsiveness plotted against photosynthetic radiation during the growth period for the annual crops that we included in our crop dataset. The relationship is positive; (b) beehive plot of mycorrhizal responsiveness of maize lines adapted to growth under temperate vs tropical conditions. In the plot, we only used the two central quartiles of the two groups of data. The differences are significant ($U = 4$, $P < 0.001$); (c) maximum latitude of the distribution of North American herbaceous plants plotted against the mycorrhizal responsiveness as assayed in a common garden experiment. Each cross describes the maximum latitude of a single species and the red line is the best fit line for the relationship with mycorrhizal responsiveness. We found few plants having a high mycorrhizal responsiveness occurring at high latitudes.



